

岩尾 忠	業績 (英文のみ)	2016年5月現在 67編
原著論文	61編 (筆頭執筆 46編、第2執筆 15編)	
総 説	1編	
コメント	3編	
症例報告	2編	

67: Iwao T, Sakai K, Ando E. Liraglutide with or without Metformin Ameliorates Liver Function and Fatty Liver in Obese Patients with Type 2 Diabetes Mellitus. *J Diabetes Metab* 2015; 6:9

66: Iwao T, Sakai K, Ando E. Relative contribution of insulin secretion and insulin sensitivity at different stage of glucose tolerance: non-obese versus obese subjects. *Internal medicine* 2014;53:383-390

65: Iwao T, Sakai K, Ando E. Seasonal fluctuations of glycated hemoglobin levels in Japanese diabetic patients; Effect of diet therapy and physical activity. *Diabetology International* 2013;4: 173-178

64: Iwao T, Sakai K, Ando E. Meaning of upper limit of normal range of post-load one-hour plasma glucose level defined by oral glucose tolerance test in Japanese subjects. *J Diabetes Invest* 2013;4: 372-375

63: Iwao T, Sakai K, Ando E. Upper limit of fasting plasma glucose level defined by oral glucose tolerance test in Japanese subjects. *Diabetes Res Clin Pract* 2013;99: 272-276

62: Iwao T, Sakai K, Sata M. Postprandial serum C-peptide is a useful parameter in the prediction of successful switching to liraglutide monotherapy from complex insulin therapy in Japanese patients with type 2 diabetes. *J Diabetes Complications*. 2013;27: 89-91

61: Sakai K, Iwao T, Oho K, Toyonaga A, Sata M. Propranolol ameliorates thrombocytopenia in patients with cirrhosis. *J Gastroenterol*. 2002;37:112-8.

60: Toyonaga A, Iwao T. Paraesophageal collaterals in endoscopic therapies for esophageal varices: good or bad? *J Gastroenterol Hepatol*. 2001;16:489-90.

- 59: Iwao T, Oho K, Nakano R, Sakai T, Sato M, Miyamoto Y, Kumamoto M, Sakai K, Sata M, Toyonaga A. High plasma cardiac natriuretic peptides associated with enhanced cyclic guanosine monophosphate production in preascitic cirrhosis. *J Hepatol.* 2000;32:426-33.
- 58: Nakano R, Iwao T, Oho K, Ono N, Sakai T, Sato M, Yamawaki M, Miyamoto Y, Sakai K, Takeda T, Tsuruta O, Sata M, Toyonaga A. Effect of transcatheter arterial chemoembolization on kidney hemodynamics and function in patients with cirrhosis and hepatocellular carcinoma. *J Hepatol.* 1999;31:340-6.
- 57: Iwao T, Oho K, Nakano R, Yamawaki M, Sakai T, Sato M, Miyamoto Y, Sakai K, Sata M, Toyonaga A. Upright posture blunts postprandial splanchnic hyperemia in patients with cirrhosis and portal hypertension. *J Gastroenterol.* 1999;34:359-65.
- 56: Toyonaga A, Iwao T. Is stab with strangulation the fastest way to kill esophageal varices? *J Gastroenterol.* 1999;34:286-7.
- 55: Toyonaga A, Iwao T, Sumino M, Takagi K, Oho K, Shigemori H, Tanikawa K. Distinctive portal venographic pattern in patients with sclerotherapy resistant oesophageal varices. *J Gastroenterol Hepatol.* 1999;11:1110-4.
- 54: Iwao T, Oho K, Nakano R, Yamawaki M, Sakai T, Sato M, Miyamoto Y, Toyonaga A, Tanikawa K. Effect of meal induced splanchnic arterial vasodilatation on renal arterial haemodynamics in normal subjects and patients with cirrhosis. *Gut.* 1998;43:843-8.
- 53: Toyonaga A, Iwao T. Portal-hypertensive gastropathy. *J Gastroenterol Hepatol.* 1998;13:865-77. Review..
- 52: Iwao T, Oho K, Sakai T, Sato M, Nakano R, Yamayaki M, Toyonaga A, Tanikawa K. Noninvasive hemodynamic measurements of superior mesenteric artery in the prediction of portal pressure response to propranolol. *J Hepatol.* 1998;28:847-55.
- 51: Nakayama M, Iwao T, Oho K, Toyonaga A, Tanikawa K. Role of extravariceal collateral channels in the development of portal-hypertensive gastropathy before and after sclerotherapy. *J Gastroenterol.* 1998;33:142-6.

- 50: Iwao T, Oho K, Sakai T, Sato M, Nakano R, Yamawaki M, Toyonaga A, Tanikawa K. Upright posture decreases esophageal varices flow velocity in patients with cirrhosis. *J Hepatol*. 1998;28:447-53.
- 49: Tayama C, Iwao T, Oho K, Toyonaga A, Tanikawa K. Effect of large fundal varices on changes in gastric mucosal hemodynamics after endoscopic variceal ligation. *Endoscopy*. 1998;30:25-31.
- 48: Sakaki M, Iwao T, Oho K, Toyonaga A, Tanikawa K. Prognostic factors in cirrhotic patients receiving long-term sclerotherapy for the first bleeding from oesophageal varices. *Eur J Gastroenterol Hepatol*. 1998;10:21-6.
- 47: Masumoto H, Toyonaga A, Oho K, Iwao T, Tanikawa K. Ligation plus low-volume sclerotherapy for high-risk esophageal varices: comparisons with ligation therapy or sclerotherapy alone. *J Gastroenterol*. 1998;33:1-5.
- 46: Nakahara K, Iwao T, Toyonaga A, Miyazaki S, Tanikawa K. Effects of glucagon on superior mesenteric artery and femoral artery haemodynamics in humans. *Eur J Gastroenterol Hepatol*. 1997;9:1233-7.
- 45: Iwao T, Oho K, Sakai T, Tayama C, Sato M, Nakano R, Yamawaki M, Toyonaga A, Tanikawa K. Splanchnic and extrasplanchnic arterial hemodynamics in patients with cirrhosis. *J Hepatol*. 1997;27:817-23.
- 44: Sakai T, Iwao T, Oho K, Toyonaga A, Tanikawa K. Influence of extravariceal collateral channel pattern on recurrence of esophageal varices after sclerotherapy. *J Gastroenterol*. 1997;32:715-9.
- 43: Nakano R, Iwao T, Oho K, Toyonaga A, Tanikawa K. Splanchnic hemodynamic pattern and liver function in patients with cirrhosis and esophageal or gastric varices. *Am J Gastroenterol*. 1997;92:2085-9.
- 42: Iwao T, Toyonaga A, Sato M, Oho K, Sakai T, Tayama C, Nakano R, Tanikawa K. Effect of posture-induced blood volume expansion on systemic and regional hemodynamics in patients with cirrhosis. *J Hepatol*. 1997;27:484-91.
- 41: Iwao T, Toyonaga A, Oho K, Sakai T, Tayama C, Masumoto H, Sato M, Nakahara K, Tanikawa K. Portal-hypertensive gastropathy develops less in patients with cirrhosis and fundal varices. *J Hepatol*. 1997;26:1235-41.

- 40: Iwao T, Toyonaga A, Oho K, Tayama C, Masumoto H, Sakai T, Sato M, Tanikawa K. Value of Doppler ultrasound parameters of portal vein and hepatic artery in the diagnosis of cirrhosis and portal hypertension. *Am J Gastroenterol.* 1997;92:1012-7.
- 39: Iwao T, Toyonaga A, Oho K, Sakai T, Tayama C, Masumoto H, Sato M, Nakahara K, Tanikawa K. Postprandial splanchnic hemodynamic response in patients with cirrhosis of the liver: evaluation with "triple-vessel" duplex US. *Radiology.* 1996;201:711-5.
- 38: Iwao T, Toyonaga A, Oho K, Shigemori H, Sakai T, Tayama C, Masumoto H, Sato M, Tanikawa K. Effect of vasopressin on esophageal varices blood flow in patients with cirrhosis: comparisons with the effects on portal vein and superior mesenteric artery blood flow. *J Hepatol.* 1996;25:491-7.
- 37: Iwao T, Toyonaga A, Ikegami M, Shigemori H, Oho K, Sumino M, Tanikawa K. Gastric mucus generation in cirrhotic patients with portal hypertension. Effects of tetraprenylacetone. *Dig Dis Sci.* 1996;41:1727-32.
- 36: Iwao T, Toyonaga A, Shigemori H, Oho K, Sakai T, Tayama C, Masumoto H, Sato M, Tanikawa K. Hepatic artery hemodynamic responsiveness to altered portal blood flow in normal and cirrhotic livers. *Radiology.* 1996;200:793-8.
- 35: Iwao T, Toyonaga A, Shigemori H, Oho K, Sumino M, Sato M, Tanikawa K. Vasopressin plus oxygen vs vasopressin alone in cirrhotic patients with portal-hypertensive gastropathy: effects on gastric mucosal haemodynamics and oxygenation. *J Gastroenterol Hepatol.* 1996;11:216-22.
- 34: Iwao T, Toyonaga A, Shigemori H, Oho K, Sumino M, Sato M, Tanikawa K. Echo-Doppler measurements of portal vein and superior mesenteric artery blood flow in humans: inter- and intra-observer short-term reproducibility. *J Gastroenterol Hepatol.* 1996;11:40-6.
- 33: Iwao T, Toyonaga A, Shigemori H, Sumino M, Oho K, Tanikawa K. Supplemental oxygen during endoscopic variceal ligation: effects on arterial oxygenation and cardiac arrhythmia. *Am J Gastroenterol.* 1995;90:2186-90.

- 32: Iwao T, Toyonaga A, Kuboyama S, Tanikawa K. Effects of omeprazole and lansoprazole on fasting and postprandial serum gastrin and serum pepsinogen A and C. *Hepatogastroenterology*. 1995;42:677-82.
- 31: Oho K, Iwao T, Sumino M, Toyonaga A, Tanikawa K. Ethanolamine oleate versus butyl cyanoacrylate for bleeding gastric varices: a nonrandomized study. *Endoscopy*. 1995;27:349-54.
- 30: Iwao T, Toyonaga A, Shigemori H, Ikegami M, Tanikawa K. Omeprazole administration does not impair gastrointestinal mucosal perfusion, oxygenation, and hexosamine generation. *Gastrointest Endosc*. 1995;41:497-501.
- 29: Iwao T, Toyonaga A, Ikegami M, Sumino M, Oho K, Sakaki M, Shigemori H, Tanikawa K, Iwao J. Effects of exercise-induced sympathoadrenergic activation on portal blood flow. *Dig Dis Sci*. 1995;40:48-51.
- 28: Iwao T, Toyonaga A, Ikegami M, Sumino M, Oho K, Sakaki M, Shigemori H, Nakayama M, Tanikawa K. Wedged hepatic venous pressure reflects portal venous pressure during vasoactive drug administration in nonalcoholic cirrhosis. *Dig Dis Sci*. 1994;39:2439-44.
- 27: Shigemori H, Iwao T, Ikegami M, Toyonaga A, Tanikawa K. Effects of propranolol on gastric mucosal perfusion and serum gastrin level in cirrhotic patients with portal hypertensive gastropathy. *Dig Dis Sci*. 1994;39:2433-8.
- 26: Toyonaga A, Iwao T, Sumino M, Oho K, Ikegami M, Sakaki M, Shigemori H, Nakayama M, Sasaki E, Tanikawa K. Portal pressure after prophylactic sclerotherapy in patients with high-risk varices. *J Hepatol*. 1994;21:515-20.
- 25: Iwao T, Toyonaga A, Ikegami M, Sumino M, Oho K, Sakaki M, Shigemori H, Nakayama M, Tanikawa K, Iwao J. Cardiovascular responsiveness after isotonic exercise in cirrhotic patients: study on sympathoadrenergic and renin-angiotensin systems. *Am J Gastroenterol*. 1994;89:1043-6.
- 24: Iwao T, Toyonaga A, Ikegami M, Shigemori H, Oho K, Sumino M, Sakaki M, Nakayama M, Nishiyama T, Minetoma T, et al. McCormack's endoscopic signs for diagnosing portal hypertension: comparison with gastroesophageal varices. *Gastrointest Endosc*. 1994;40:470-3.

- 23: Iwao T, Toyonaga A, Ikegami M, Sumino M, Oho K, Shigemori H, Sakaki M, Nakayama M, Tanikawa K, Iwao J. Portal vein hemodynamics in cirrhotic patients with portal hypertensive gastropathy: an echo-Doppler study. *Hepatogastroenterology*. 1994;41:230-4.
- 22: Iwao T, Toyonaga A, Harada H, Harada K, Ban S, Minetoma T, Sumino M, Ikegami M, Tanikawa K. Arterial oxygen desaturation during non-sedated diagnostic upper gastrointestinal endoscopy in patients with cirrhosis. *Gastrointest Endosc*. 1994;40:281-4.
- 21: Iwao T, Toyonaga A, Harada H, Harada K, Ban S, Ikegami M, Tanikawa K. Arterial oxygen desaturation during non-sedated diagnostic upper gastrointestinal endoscopy. *Gastrointest Endosc*. 1994;40:277-80.
- 20: Ikegami M, Toyonaga A, Iwao T, Tarikawa K. Chronic lymphocytic leukemia with esophageal varices. *Dig Dis Sci*. 1993;38:2142-3.
- 19: Iwao T, Toyonaga A, Ikegami M, Oho K, Sumino M, Sakaki M, Shigemori H, Nakayama M, Sasaki E, Tanikawa K. Effects of vasopressin and nicardipine on hemodynamics and liver function in patients with cirrhosis: comparison with vasopressin alone. *J Hepatol*. 1993;19:345-52.
- 18: Iwao T, Toyonaga A. "Passive" gastric mucosal congestion in patients with gastropathy. *Dig Dis Sci*. 1993;38:1563-4.
- 17: Iwao T, Toyonaga A, Ikegami M. Gastric mucosal hemodynamics in patients with portal hypertensive gastropathy. *Gastroenterology*. 1993;105:309-10.
- 16: Iwao T, Toyonaga A, Ikegami M, Oho K, Sumino M, Harada H, Sakaki M, Shigemori H, Aoki T, Tanikawa K. Reduced gastric mucosal blood flow in patients with portal-hypertensive gastropathy. *Hepatology*. 1993;18:36-40.
- 15: Iwao T, Toyonaga A, Ikegami M, Sumino M, Oho K, Sakaki M, Shigemori H, Tanikawa K, Iwao J. Reduced portosystemic hemodynamic responsiveness after orthostasis in patients with cirrhosis. *Dig Dis Sci*. 1993;38:1251-8.
- 14: Iwao T, Toyonaga A, Kim M, Sumino M, Ikegami M, Oho K, Ikeda H, Majima Y, Sasaki E, Tanikawa K. Prevalence of gastroesophageal varices supplied by intrahepatic portal branch. *Intern Med*. 1993;32:462-4.

- 13: Iwao T, Toyonaga A, Ikegami M, Tanikawa K. Portal and gastric mucosal hemodynamics after sclerotherapy. *Gastrointest Endosc*. 1993;39:466.
- 12: Iwao T, Toyonaga A, Ikegami M, Oho K, Sumino M, Sakaki M, Shigemori H, Harada H, Sasaki E, Tanikawa K. Gastric mucosal blood flow after smoking in healthy human beings assessed by laser Doppler flowmetry. *Gastrointest Endosc*. 1993;39:400-3.
- 11: Iwao T, Toyonaga A, Ikegami M, Oho K, Sumino M, Sakaki M, Harada H, Tanikawa K. Observer agreement and variability in measuring gastric mucosal blood flow by laser Doppler flowmetry in humans. *Endoscopy*. 1993;25:274-7.
- 10: Iwao T, Toyonaga A, Ikegami M, Sumino M, Oho K, Sakaki M, Nishizono M, Tanikawa K. Portohepatic pressures, hepatic function, and blood gases in the combination of nitroglycerin and vasopressin: search for additive effects in cirrhotic portal hypertension. *Am J Gastroenterol*. 1992;87:719-24.
- 9: Iwao T, Toyonaga A, Sumino M, Takagi K, Oho K, Nishizono M, Ohkubo K, Inoue R, Sasaki E, Tanikawa K. Portal hypertensive gastropathy in patients with cirrhosis. *Gastroenterology*. 1992;102:2060-5.
- 8: Iwao T, Toyonaga A, Sumino M, Oho K, Sakaki M, Ikegami M, Nishizono M, Tanikawa K. Lack of hepatic benefit by oxygen inhalation during vasopressin infusion in patients with cirrhosis. *J Gastroenterol Hepatol*. 1992;7:249-52.
- 7: Iwao T, Toyonaga A, Sumino M, Takagi K, Oho K, Ohkubo K, Inoue R, Tanikawa K. Effects of vaso-active agents on hepatic function and blood gases in patients with cirrhosis: a study of vasopressin and nitroglycerin. *J Gastroenterol Hepatol*. 1992;7:148-53.
- 6: Iwao T, Toyonaga A, Ikegami M, Sugawara H, Sumino M, Takagi K, Oho K, Sakaki M, Ohkubo K, Inoue R, et al. Nicardipine infusion improved hepatic function but failed to reduce hepatic venous pressure gradient in patients with cirrhosis. *Am J Gastroenterol*. 1992;87:326-31.
- 5: Yamashita F, Iwao T, Torimura T, Tanaka M, Hirai K, Abe M, Toyonaga A, Sugihara S, Kojiro M, Tanikawa K. Sclerosing hepatocellular carcinoma with hypercalcemia--a case report. *Kurume Med J*. 1992;39:113-6.

- 4: Iwao T, Toyonaga A, Sumino M, Takagi K, Ohkubo K, Inoue R, Tanikawa K. Effect of intraportal nitroglycerin on portosystemic hemodynamics in patients with cirrhosis: comparison with i.v. route. *J Clin Gastroenterol*. 1991;13:424-31.
- 3: Iwao T, Toyonaga A, Sumino M, Takagi K, Ohkubo K, Inoue R, Tanikawa K. Hemodynamic study during transdermal application of nitroglycerin tape in patients with cirrhosis. *Hepatology*. 1991;13:124-8.
- 2: Iwao T, Toyonaga A, Tanikawa K. Gastric red spots in patients with cirrhosis: subclinical condition of gastric mucosal hemorrhage? *Gastroenterol Jpn*. 1990;25:685-92.
- 1: Toyonaga A, Iwao T, Shimotsuura Y, Tanikawa K. Endoscopic, histologic and haemodynamic studies on portal hypertensive gastric mucosa. *J Gastroenterol Hepatol*. 1989;4 Suppl 1:132-5.